

HUPD-9316
YNU-HEPTh-93-102
November 1993

The Mass of the Lightest Supersymmetric Higgs Boson beyond the Leading Logarithm Approximation

JIRO KODAIRA* AND YOSHIAKI YASUI [†]

*Dept. of Physics, Hiroshima University
Higashi-Hiroshima 724, JAPAN*

KEN SASAKI[‡]

*Dept. of Physics, Yokohama National University
Yokohama 240, JAPAN*

Abstract

We examine the radiative corrections to the mass of the lightest Higgs boson in the minimal supersymmetric extension of the standard model. We use the renormalization-group improved effective potential which includes the next-to-leading-order contributions. We find that, contrary to the result of Espinosa and Quirós, the higher-order corrections to the lightest Higgs boson mass are non-negligible, adding 3 – 11 GeV (3 – 9 GeV) to the result in the leading logarithm approximation for the range of top quark mass $100\text{GeV} < m_t < 200\text{GeV}$ and for the supersymmetric breaking scale $M_{\text{SUSY}} = 1\text{TeV}$ ($M_{\text{SUSY}} = 10\text{TeV}$). Also we find that our result is stable under the change of the renormalization parameter t .

*Work partially supported by the Monbusho Grant-in-Aid for Scientific Research (C) No. 05640351.

[†]Fellow of the Japan Society for the Promotion of Science for Japanese Junior Scientist. Work partially supported by the Monbusho Grant-in-Aid for Scientific Research No. 050076.

[‡]e-mail address: a121004@c1.ed.ynu.ac.jp

Although the standard model (SM) is highly successful and in excellent agreement with all the measurements at the present energies, it is widely believed that SM is not the final theory for the world of elementary particles. The minimal supersymmetric extension of the standard model (MSSM) is one of the most promising candidates beyond the SM. The MSSM possesses in its physical spectrum three neutral and two charged Higgs bosons, and there exists a tree-level relation which implies that at least one neutral Higgs boson is lighter than the Z^0 mass (M_Z). Radiative corrections to the masses of these Higgs bosons have been calculated by several groups [1] - [14], who found that they are quite significant, depending strongly on the top quark mass and the scale of supersymmetry breaking (M_{SUSY}) or the squark masses. All the above works except ref. [6] considered the one-loop radiative corrections to Higgs boson masses.

Now that the one-loop corrections have been found to be significantly large, it is quite natural to ask next how large the higher-order corrections would be. Indeed Espinosa and Quirós [6] have analyzed the “two-loop” radiative corrections to the mass of the lightest Higgs boson in the minimal and non-minimal (including a gauge singlet) supersymmetric standard model. They used the effective potential (EP) in the leading logarithm approximation and examined the evolution of the Higgs quartic coupling λ by the renormalization group (RG) techniques with the one- and two-loop β functions. They found that the “two-loop” correction is negative and stays within a few percent even in cases where the one-loop correction is larger than the tree-level mass.

Recently there appeared interesting papers [15] - [17] which discussed about the improvement of EP by using the renormalization group equation (RGE) . It was shown there that to improve EP which satisfies the RGE with up to the two-loop β functions and anomalous dimension γ , one should include the one-loop-level potential with the running parameters into the solution. In this respect, the work of Espinosa and Quirós [6] seems unsatisfactory: they used the EP in the leading logarithm approximation, which is the tree-level potential with the running parameters, and they made use of the two-loop β functions only to determine the evolution of these parameters. In this paper we reanalyze the mass of the lightest

Higgs boson (m_ϕ) in the MSSM using the EP improved by RGE up to the next-to-leading order. We find that new terms which were not considered by Espinosa and Quirós give non-negligible contributions to the m_ϕ . We also find that the predicted values of m_ϕ are stable under the change of the renormalization parameter t when we use the RGE-improved EP which includes the next-to-leading-order contributions.

Two basic assumptions were made in their analysis of the lightest Higgs boson mass [6]: (a) all supersymmetric (SUSY) partners of the SM particles have masses of the order of the supersymmetry breaking scale M_{SUSY} ; (b) one linear combination H of the two Higgs boson doublets, $H_1 = (H_1^0, H_1^-)^T$ and $H_2 = (H_2^+, H_2^0)^T$,

$$H = H_1 \cos\beta + i\tau_2 H_2^* \sin\beta \quad (1)$$

is light, while the other linear combination, which is orthogonal to the former one, is as heavy as the SUSY partners. Under these assumptions, it is clear that the effective theory below the scale M_{SUSY} is the usual SM with one light Higgs doublet H . Throughout the following analyses, we will make the same assumptions (a) and (b). The tree-level Higgs potential below M_{SUSY} is then given by

$$V_{\text{tree}} = -m^2 |H|^2 + \frac{1}{6} \lambda |H|^4, \quad (2)$$

where

$$\frac{1}{3} \lambda = \frac{1}{4} (g_1^2 + g_2^2) \cos^2 2\beta, \quad (3)$$

and g_1 and g_2 are the gauge coupling constants of $U(1)_Y$ and $SU(2)_L$, respectively.

When the neutral component of H acquires a vacuum expectation value $\langle H^0 \rangle = v/\sqrt{2}$, the above tree-level potential (2) gives the physical Higgs $\phi \equiv (\text{Re} H^0 - v)/\sqrt{2}$ (which corresponds to the lightest Higgs boson under the assumptions (a) and (b)) a squared mass

$$m_\phi^2 = \frac{1}{3} \lambda v^2. \quad (4)$$

Also at the symmetry breaking, the top quark and the Z^0 , W^\pm gauge bosons gain masses which are given by

$$m_t = h_t v / \sqrt{2}, \quad M_Z^2 = \frac{1}{4} (g_1^2 + g_2^2) v^2, \quad M_W^2 = \frac{1}{4} g_2^2 v^2, \quad (5)$$

where h_t is the Yukawa coupling of ϕ to the top quark. The tree-level relation $m_\phi^2 = M_Z^2 \cos^2 2\beta$ follows from Eqs. (3) - (5).

The (RGE-unimproved) EP of the SM up to the one-loop level is given by

$$V_1 = V_{(0)} + V_{(1)}, \quad (6)$$

$$V_{(0)} = -\frac{1}{2}m^2\varphi_c^2 + \frac{1}{24}\lambda\varphi_c^4, \quad (7)$$

$$V_{(1)} = -\frac{3}{64\pi^2}(h_t^2\varphi_c^2)^2\left(\ln\frac{h_t^2\varphi_c^2}{2\mu^2} - \frac{3}{2}\right) + \dots, \quad (8)$$

where φ_c is the classical field corresponding to the physical Higgs boson ϕ , and all the Yukawa couplings of ϕ to quarks and leptons except the top quark are neglected. The calculation is performed in the Landau gauge and in the \overline{MS} scheme to obtain the one-loop result $V_{(1)}$, and μ is the renormalization scale. The ellipses in Eq. (8) represent contributions of the gauge bosons and the would-be-Goldstone bosons. Throughout this paper we use the Landau gauge which is the most convenient for our purpose [18] and the \overline{MS} scheme.

Now we improve the EP by using the RGE. It was recently emphasized by the authors of Refs. [15] - [17] that in the \overline{MS} scheme the EP $V(\varphi_c)$ fails to satisfy the usual (homogeneous) RGE unless $V(0)$, a contribution to the “vacuum energy”, is suitably dealt with. When we use the RGE-improved EP in the leading order and obtain m_ϕ , the consideration of $V(0)$ term is unnecessary. However, as we shall see below, if we improve the EP by RGE up to the next-to-leading order, $V(0)$ becomes relevant to us and we must take its presence into account. Thus with an appropriate φ_c -independent term being added, the new EP $V(\varphi_c)$ satisfies the following RGE of the usual form:

$$(\mathcal{D} - \gamma_\phi\varphi_c\frac{\partial}{\partial\varphi_c})V(\varphi_c, X_i, \mu) = 0 \quad (9)$$

with

$$\mathcal{D} = \mu\frac{\partial}{\partial\mu} + \beta_{X_i}\frac{\partial}{\partial X_i}, \quad (10)$$

where $X_i = \lambda, h_t, m^2, g_3, g_2, g_1$ and g_3 is the gauge coupling constant of $SU(3)_C$. The solution is easily found by the method of characteristics and we get

$$V(\varphi_c, X_i, \mu) = V(\varphi_c(t), X_i(t), \mu(t)), \quad (11)$$

where $X_i(t)$ are running couplings and running mass which are determined by the equations

$$\frac{dX_i(t)}{dt} = \beta_{X_i}(X_j(t)), \quad X_i, X_j = \lambda, h_t, m^2, g_3, g_2, g_1 \quad (12)$$

with the boundary conditions $X_i(0) = X_i$, and

$$\begin{aligned} \xi(t) &= \exp \left\{ - \int_0^t \gamma_\phi(t') dt' \right\}, \\ \varphi_c(t) &= \varphi_c \xi(t), \\ \mu(t) &= \mu e^t. \end{aligned} \quad (13)$$

Then using the result of tree- and one-loop-level EP of Eqs.(7) and (8), we obtain the RGE-improved V as follows:

$$\begin{aligned} V &= \Omega(X_i(t), \mu(t)) \\ &\quad + V_{(0)}(\varphi_c(t), X_i(t)) \\ &\quad + V_{(1)}(\varphi_c(t), X_i(t), \mu(t)) + \cdots, \end{aligned} \quad (14)$$

where Ω is the φ_c -independent term which is added for V to satisfy a RGE of the usual homogeneous form, and the ellipses represent the higher-loop contributions.

For later convenience, let us expand the RGE coefficient functions β_{X_i} and γ_ϕ by the number of loops as follows:

$$\begin{aligned} \beta_{X_i} &= \hbar \beta_{X_i}^{(1)} + \hbar^2 \beta_{X_i}^{(2)} + \cdots, \quad X_i = \lambda, h_t, m^2, g_3, g_2, g_1 \\ \gamma_\phi &= \hbar \gamma_\phi^{(1)} + \hbar^2 \gamma_\phi^{(2)} + \cdots, \end{aligned} \quad (15)$$

where we have introduced the Plank's constant \hbar so that the power of \hbar counts the number of loops and $\beta_{X_i}^{(n)}$ and $\gamma_\phi^{(n)}$ are the n -loop contribution to β_{X_i} and γ_ϕ , respectively. Similarly V has the loop expansion

$$V = \Omega + V_{(0)}(\varphi_c(t)) + \hbar V_{(1)}(\varphi_c(t)) + \cdots, \quad (16)$$

and we have denoted $\Omega(X_i(t), \mu(t))$ and $V_{(n)}(\varphi_c(t), X_i(t), \mu(t))$ as Ω and $V_{(n)}(\varphi_c(t))$, for short, respectively.

Inserting Eq.(16) into Eq.(9) and picking the terms up to of the order \hbar , we find

$$\mathcal{D}\Omega + \hbar \left\{ \beta_{X_i}^{(1)} \frac{\partial V_{(0)}}{\partial X_i} - \gamma_\phi^{(1)} \varphi_c \frac{\partial V_{(0)}}{\partial \varphi_c} + \frac{3}{32\pi^2} h_t^4 \varphi_c^4 \right\} = 0, \quad (17)$$

where the last term in the parentheses arises from $\mu \partial V_{(1)} / \partial \mu$ when we use the expression of $V_{(1)}$ in Eq.(8) and neglect the contributions of the gauge bosons and the would-be-Goldstone bosons to $V_{(1)}$. If we further set $\varphi_c = 0$ in the above equation, we obtain

$$\mathcal{D}\Omega + \hbar \mu \frac{\partial V_{(1)}(\varphi_c = 0)}{\partial \mu} = 0, \quad (18)$$

from which Ω can be determined to the leading order. For the later discussion, however, we do not need the specific form of Ω , and we will see below that the knowledge of Eq.(17) is sufficient for our purpose.

We will now analyze the mass of the lightest Higgs boson using the RGE-improved V . At first, let us consider the boundary conditions for coupling constants. Under our basic assumptions explained before, the relation between the quartic coupling constant λ and the gauge coupling constants g_1 and g_2 given in Eq.(3) should be satisfied at the scale M_{SUSY} , that is,

$$\frac{1}{3}\lambda(M_{\text{SUSY}}) = \frac{1}{4}(g_1^2(M_{\text{SUSY}}) + g_2^2(M_{\text{SUSY}}))\cos^2 2\beta. \quad (19)$$

So we choose the renormalization scale μ to be M_{SUSY} and take the parameter t as $t = \ln(\varphi_c/M_{\text{SUSY}})$. Then we find that $\mu(t) = \varphi_c$ and the RGE-improved V is given by

$$V = \Omega + V_{(0)}(\varphi_c(t)) + \hbar V_{(1)}(\varphi_c(t)) + \mathcal{O}(\hbar^2), \quad (20)$$

with

$$V_{(1)}(\varphi_c(t)) = -\frac{3}{64\pi^2} h_t^4(t) \varphi_c^4(t) \left[\ln \frac{h_t^2(t) \xi^2(t)}{2} - \frac{3}{2} \right], \quad (21)$$

where in $V_{(1)}(\varphi_c(t))$ we only include the top-quark contribution to the one-loop EP, because, with its very large Yukawa coupling, the contribution of top-quark is dominant over those from the gauge bosons and the would-be Goldstone bosons. We will obtain the lightest Higgs boson mass by evaluating $\partial^2 V / \partial \varphi_c(t)^2$ at $\varphi_c(t_v) = v = (\sqrt{2}G_F)^{-\frac{1}{2}} = 246\text{GeV}$ under the minimum condition $\partial V / \partial \varphi_c(t) = 0$ at $\varphi_c(t_v) = v$. The value t_v is determined by the equation $\varphi_c(t_v) = v$, which, with help of Eq.(13), is transformed into

$$t_v = \ln \frac{v}{M_{\text{SUSY}}} + \int_0^{t_v} \gamma_\phi(t') dt'. \quad (22)$$

It is noted that we differentiate V not by φ_c but by the renormalized field $\varphi_c(t)$ and also that we evaluate the differentials at the point of $\varphi_c(t_v) = v$ and not at $\varphi_c = v$.

Since $X_i(t)$, $\varphi_c(t)$, and $\mu(t)$ are functions of t , we find

$$\begin{aligned}\frac{\partial X_i(t)}{\partial \varphi_c(t)} &= \hbar \beta_{X_i}^{(1)}(t) \frac{1}{\varphi_c(t)} + \mathcal{O}(\hbar^2) \\ \frac{\partial \mu(t)}{\partial \varphi_c(t)} &= \frac{\mu(t)}{\varphi_c(t)} + \mathcal{O}(\hbar).\end{aligned}\tag{23}$$

Thus we obtain

$$\begin{aligned}\frac{\partial \Omega}{\partial \varphi_c(t)} &= \frac{1}{\varphi_c(t)} \left\{ \mu(t) \frac{\partial \Omega}{\partial \mu(t)} + \hbar \beta_{X_i}^{(1)}(t) \frac{\partial \Omega}{\partial X_i(t)} \right\} + \mathcal{O}(\hbar^2), \\ \frac{\partial V_{(0)}(\varphi_c(t))}{\partial \varphi_c(t)} &= V'_{(0)}(\varphi_c(t)) + \hbar \frac{\beta_{X_i}^{(1)}(t)}{\varphi_c(t)} \frac{\partial V_{(0)}(\varphi_c(t))}{\partial X_i(t)} + \mathcal{O}(\hbar^2), \\ \frac{\hbar \partial V_{(1)}(\varphi_c(t))}{\partial \varphi_c(t)} &= -\hbar \frac{3}{16\pi^2} h_t^4(t) \varphi_c^3(t) \left[\ln \frac{h_t^2(t) \xi^2(t)}{2} - \frac{3}{2} \right] + \mathcal{O}(\hbar^2).\end{aligned}\tag{24}$$

It is now straightforward to evaluate $\partial V / \partial \varphi_c(t)$ from Eq.(24). Using the relation (17), we eliminate $\mathcal{D}\Omega$ term and find

$$\begin{aligned}\frac{\partial V}{\partial \varphi_c(t)} &= (1 + \hbar \gamma_\phi^{(1)}(t)) \left\{ -m^2(t) \varphi_c(t) + \frac{1}{6} \lambda(t) \varphi_c^3(t) \right\} \\ &\quad + \hbar \left\{ -\frac{3}{16\pi^2} h_t^4(t) \varphi_c^3(t) \left[\ln \frac{h_t^2(t) \xi^2(t)}{2} - 1 \right] \right\} \\ &\quad + \mathcal{O}(\hbar^2).\end{aligned}\tag{25}$$

Further differentiation of $\partial V / \partial \varphi_c(t)$ by $\varphi_c(t)$ gives

$$\begin{aligned}\frac{\partial^2 V}{\partial \varphi_c^2(t)} &= (1 + \hbar \gamma_\phi^{(1)}(t)) \left\{ -m^2(t) + \frac{1}{2} \lambda(t) \varphi_c^2(t) \right\} \\ &\quad + \hbar \left\{ -\beta_{m^2}^{(1)}(t) + \frac{1}{6} \beta_\lambda^{(1)}(t) \varphi_c^2(t) \right\} \\ &\quad + \hbar \left\{ -\frac{9}{16\pi^2} h_t^4(t) \varphi_c^2(t) \left[\ln \frac{h_t^2(t) \xi^2(t)}{2} - 1 \right] \right\} \\ &\quad + \mathcal{O}(\hbar^2).\end{aligned}\tag{26}$$

Using the minimum condition $\partial V/\partial\varphi_c(t) = 0$ at $\varphi_c(t_v) = v$, we finally obtain for the lightest Higgs mass in the next-to-leading logarithm approximation

$$\begin{aligned}
m_{\phi(2\text{-loop})}^2 &= \left. \frac{\partial^2 V}{\partial\varphi_c^2(t)} \right|_{\varphi_c(t_v)=v} \\
&= \frac{1}{3}\lambda(t_v)v^2 \\
&\quad + \hbar v^2 \left\{ \frac{1}{6}\beta_\lambda^{(1)}(t_v) - \frac{1}{6}\lambda(t_v) \left[\frac{\beta_{m^2}^{(1)}(t_v)}{m^2(t_v)} - 2\gamma_\phi^{(1)}(t_v) \right] \right. \\
&\quad \left. - \frac{3}{8\pi^2}h_t(t_v)^4 \left[\ln \frac{h_t^2(t_v)\xi^2(t_v)}{2} - 1 \right] \right\} \\
&\quad + \mathcal{O}(\hbar^2).
\end{aligned} \tag{27}$$

The first term in Eq.(27) gives

$$m_{\phi(1\text{-loop})}^2 = (1/3)\lambda(t_v)v^2, \tag{28}$$

which is the result given by Refs. [1] [4] except that the authors of Ref. [1] have evaluated the running coupling $\lambda(t)$ at $t_{m_\phi} = \ln(m_\phi/M_{\text{SUSY}})$. As far as the leading logarithm approximation is concerned, the terms of order \hbar in Eq.(27) are neglected as the higher-order effects. An arbitrariness coming from different choice of the parameter t also falls in the higher-order corrections although it has no small effect on the predicted values numerically. Espinosa and Quirós have employed the “one-loop” formula $m_\phi^2 = (1/3)\lambda(t)v^2$, which is correct only in the leading logarithm approximation, and computed $\lambda(t)$ at $t_{m_\phi} = \ln(m_\phi/M_{\text{SUSY}})$ using RGE with up to two-loop β functions [6]. However, if we make use of the two-loop RGE coefficient functions for the running parameters and evaluate the lightest Higgs mass m_ϕ^2 , we should take into account the order- \hbar -terms in Eq.(27) which also collect the next-to-leading logarithmic contributions.

The one- and two-loop β functions and anomalous dimension γ_ϕ for the SM, which we will use in this analysis, read as follows [17]. We define the constant A as $A \equiv 16\pi^2$.

For the Higgs quartic coupling λ :

$$A\beta_\lambda^{(1)} = 4\lambda^2 + 12\lambda h_t^2 - 36h_t^4 - 3\lambda(3g_2^2 + g_1^2) + \frac{27}{4}g_2^4 + \frac{9}{2}g_2^2g_1^2 + \frac{9}{4}g_1^4,$$

$$\begin{aligned}
A^2\beta_\lambda^{(2)} &= -\frac{26}{3}\lambda^3 - 24\lambda^2 h_t^2 + 6\lambda^2(3g_2^2 + g_1^2) \\
&\quad + \lambda\{-3h_t^4 + h_t^2(80g_3^2 + \frac{45}{2}g_2^2 + \frac{85}{6}g_1^2) - \frac{73}{8}g_2^4 + \frac{39}{4}g_2^2g_1^2 + \frac{629}{24}g_1^4\} \\
&\quad + 180h_t^6 - h_t^4(192g_3^2 + 16g_1^2) + h_t^2(-\frac{27}{2}g_2^4 + 63g_2^2g_1^2 - \frac{57}{2}g_1^4) \\
&\quad + \frac{915}{8}g_2^6 - \frac{289}{8}g_2^4g_1^2 - \frac{559}{8}g_2^2g_1^4 - \frac{379}{8}g_1^6.
\end{aligned} \tag{29}$$

For the top-quark Yukawa coupling h_t :

$$\begin{aligned}
A\beta_{h_t}^{(1)} &= \frac{9}{2}h_t^3 - h_t(8g_3^2 + \frac{9}{4}g_2^2 + \frac{17}{12}g_1^2), \\
A^2\beta_{h_t}^{(2)} &= h_t\{-12h_t^4 - 2\lambda h_t^2 + h_t^2(36g_3^2 + \frac{225}{16}g_2^2 + \frac{131}{16}g_1^2) \\
&\quad + \frac{1}{6}\lambda^2 - 108g_3^4 + 9g_3^2g_2^2 + \frac{19}{9}g_3^2g_1^2 - \frac{23}{4}g_2^4 - \frac{3}{4}g_2^2g_1^2 + \frac{1187}{216}g_1^4\}.
\end{aligned} \tag{30}$$

For the gauge couplings g_3 , g_2 , and g_1 :

$$\begin{aligned}
A\beta_{g_3}^{(1)} &= -7g_3^3, \\
A^2\beta_{g_3}^{(2)} &= g_3^3(-2h_t^2 - 26g_3^2 + \frac{9}{2}g_2^2 + \frac{11}{6}g_1^2), \\
A\beta_{g_2}^{(1)} &= -\frac{19}{6}g_2^3, \\
A^2\beta_{g_2}^{(2)} &= g_2^3(-\frac{3}{2}h_t^2 + 12g_3^2 + \frac{35}{6}g_2^2 + \frac{3}{2}g_1^2), \\
A\beta_{g_1}^{(1)} &= \frac{41}{6}g_1^3, \\
A^2\beta_{g_1}^{(2)} &= g_1^3(-\frac{17}{6}h_t^2 + \frac{44}{3}g_3^2 + \frac{9}{2}g_2^2 + \frac{199}{18}g_1^2).
\end{aligned} \tag{31}$$

For the mass parameter m^2 :

$$\begin{aligned}
A\beta_{m^2}^{(1)} &= m^2(2\lambda + 6h_t^2 - \frac{9}{2}g_2^2 - \frac{3}{2}g_1^2), \\
A^2\beta_{m^2}^{(2)} &= m^2\{-\frac{5}{3}\lambda^2 - 12\lambda h_t^2 + 4\lambda(3g_2^2 + g_1^2) - \frac{27}{2}h_t^4 \\
&\quad + h_t^2(40g_3^2 + \frac{45}{4}g_2^2 + \frac{85}{12}g_1^2) - \frac{145}{16}g_2^4 + \frac{15}{8}g_2^2g_1^2 + \frac{157}{48}g_1^4\}.
\end{aligned} \tag{32}$$

For the anomalous dimension γ_ϕ :

$$A\gamma_\phi^{(1)} = 3h_t^2 - \frac{9}{4}g_2^2 - \frac{3}{4}g_1^2,$$

$$\begin{aligned}
A^2\gamma_\phi^{(2)} &= \frac{1}{6}\lambda^2 - \frac{27}{4}h_t^4 + h_t^2(20g_3^2 + \frac{45}{8}g_2^2 + \frac{85}{24}g_1^2) \\
&\quad - \frac{271}{32}g_2^4 + \frac{9}{16}g_2^2g_1^2 + \frac{431}{96}g_1^4.
\end{aligned} \tag{33}$$

When we substitute the expressions of the RGE coefficient functions $\beta_\lambda^{(1)}$, $\beta_{m^2}^{(1)}$, and $\gamma_\phi^{(1)}$ into Eq.(27), we find that many terms cancel out and we obtain a rather simple expression

$$\begin{aligned}
m_{\phi(2\text{-loop})}^2 &= \frac{1}{3}\lambda v^2 \\
&\quad + \hbar v^2 \frac{1}{16\pi^2} \left\{ \frac{1}{3}\lambda^2 + 2\lambda h_t^2 - \frac{1}{2}\lambda(3g_2^2 + g_1^2) \right. \\
&\quad \left. + \frac{9}{8}g_2^4 + \frac{3}{4}g_2^2g_1^2 + \frac{3}{8}g_1^4 - 6h_t^4 \ln \frac{h_t^2 \xi^2}{2} \right\}
\end{aligned} \tag{34}$$

where it is understood that all the running parameters are the ones evaluated at $t = t_v$. It is interesting to note, in particular, that a h_t^4 term in Eq.(27), which comes from the one-loop EP of Eq.(8), cancels with another h_t^4 term in $\beta_\lambda^{(1)}$ of Eq.(29) and, in consequence, there appears only one h_t^4 term of the form $h_t^4 \ln(h_t^2 \xi^2/2)$ in the order- \hbar contributions in Eq.(34). This makes the higher-order corrections to $m_{\phi(2\text{-loop})}^2$ to be a milder one. From the expression of Eq.(34) we expect that the next-to-leading-order corrections give a positive contribution to m_ϕ^2 , which will be shown numerically to be true below.

To evaluate $m_{\phi(2\text{-loop})}$ of Eq.(34) numerically, we choose the initial conditions for the gauge couplings $\alpha_i \equiv g_i^2/4\pi$ (i=3,2,1) at the scale $M_Z = 91.2\text{GeV}$ to be

$$\alpha_3(M_Z) = 0.115, \quad \alpha_2(M_Z) = 0.0336, \quad \alpha_1(M_Z) = 0.0102 \tag{35}$$

which are consistent with present experimental constraints [19] - [21], and define the Yukawa coupling of the top quark at the scale of its mass m_t as

$$h_t(m_t) = \sqrt{2}m_t/v \quad \text{with} \quad v = 246\text{GeV}. \tag{36}$$

The fact that the two-loop $\beta_\lambda^{(2)}$, $\beta_{h_t}^{(2)}$, and $\beta_{g_i}^{(2)}$ (i=3,2,1) are functions of the couplings λ , h_t , and g_i casts Eq.(12) into a very complicated system of coupled differential equations. For given values of m_t and M_{SUSY} , we first solve the system (12) and

(22) with β_{X_i} and γ_ϕ given in Eqs.(29) - (33) together with the initial conditions Eqs.(35) and (36), and we obtain the appropriate t_v and $\lambda(t_v)$ so that λ satisfies the boundary condition, Eq.(19), when it evolves from t_v to $t = 0$. At the same time when we find the appropriate t_v and $\lambda(t_v)$, we gain all the information on the parameters which appear in the r.h.s. of Eq.(34). This is how we calculate $m_{\phi(2\text{-loop})}$ for given values of m_t and M_{SUSY} .

In Fig.1 we plot $m_{\phi(2\text{-loop})}$ as a function of m_t for $M_{\text{SUSY}} = 1\text{ TeV}$, $\cos^2 2\beta = 1$ and $\cos^2 2\beta = 0$ along with $m_{\phi(1\text{-loop})}$ in the leading logarithm approximation. Fig.2 shows the case for $M_{\text{SUSY}} = 10\text{ TeV}$. Since it is suggested that m_t is not too excessively large in MSSM [22], we have studied $m_{\phi(2\text{-loop})}$ for m_t from 100GeV to 200GeV. The curves for $\cos^2 2\beta = 1$ ($\cos^2 2\beta = 0$) can be considered as upper (lower) bounds for the lightest Higgs boson mass in the MSSM. From Fig.1 and 2 we observe that the next-to-leading-order effects are non-negligible. They add 3 – 11 GeV (3 – 9 GeV) to the result in the leading logarithm approximation for the range of top quark mass $100\text{GeV} < m_t < 200\text{GeV}$ and for $M_{\text{SUSY}} = 1\text{TeV}$ ($M_{\text{SUSY}} = 10\text{TeV}$). These rather large corrections come from the order- \hbar -terms of Eq.(34), since without those terms we could indeed recover the result of Ref. [6], namely, the higher-order corrections being negative and negligible for the considered range of parameters. Contrary to the conclusion of Espinosa and Quirós, our result shows that the higher-order corrections turn out to be positive and non-negligible when top quark is very heavy.

It is to be noted that our result in the leading logarithm approximation differs numerically from those of Refs. [1] [6] because we evaluated the running coupling $\lambda(t)$ at t_v given by Eq.(22), instead of at $t_{m_\phi} = \ln(m_\phi/M_{\text{SUSY}})$. In other words, we have made a different choice of t from the ones made in the above references. In the case of $\cos^2 2\beta = 1$, for example, our predicted values for $m_{\phi(1\text{-loop})}$ are smaller than those calculated in Refs. [1] [6] by 0 – 6GeV (0 – 10GeV) for the range $100\text{GeV} < m_t < 200\text{GeV}$ and for $M_{\text{SUSY}} = 1\text{TeV}$ ($M_{\text{SUSY}} = 10\text{TeV}$). In the leading logarithm approximation, the predicted values for the lightest Higgs boson mass are rather sensitive to the choice of the renormalization parameter t . However, in the next-to-leading logarithm approximation, it is not possible to change the

definition of t without modifying the $\mathcal{O}(\hbar)$ terms of EP. In consequence, the result is stable under the change of t . This is a well-known issue which often arises in the renormalization group approach, and will be discussed in more detail in the second comment below.

Fig.1 shows that the next-to-leading-order corrections are large for $\cos^2 2\beta = 0$ especially when $M_{\text{SUSY}} = 1\text{TeV}$. In the case of $\cos^2 2\beta = 0$, the boundary condition for $\lambda(t)$ at M_{SUSY} is $\lambda(t=0) = 0$. When $M_{\text{SUSY}} = 1\text{TeV}$, the “evolution time” is not long enough for $\lambda(t)$ to grow from the initial value $\lambda = 0$, and thus a term of the form $-6h_t^4 \ln(h_t^2 \xi^2/2)$ and gauge-coupling-constant terms in the order- \hbar contributions in Eq.(34) give relatively large corrections compared with the leading $(1/3)\lambda(t_v)v^2$.

A few comments are in order. The first comment concerns the definition of mass. The “mass” we have calculated is not the on-shell mass. It might be necessary to consider the correction coming from the wave function renormalization in order to make a realistic prediction for the Higgs mass. However, this effect is expected to be small [2] [23].

Secondly, we have taken the parameter t as $t = \ln(\varphi_c/M_{\text{SUSY}})$ to derive Eq.(27). But the physics should not depend on the choice of t . For example, as was stressed in Refs. [16] [17], the “natural choice” of t may be given by the equation

$$2\mu^2(t) = 2\mu^2 e^{2t} = h_t^2(t)\varphi_c^2(t). \quad (37)$$

Then the RGE-improved EP which we deal with will be

$$\begin{aligned} \tilde{V} = & \Omega(X_i(t), \mu(t)) - \frac{1}{2}m^2(t)\varphi_c^2(t) + \frac{1}{24}\lambda(t)\varphi_c^4(t) \\ & + \hbar \left\{ \frac{9}{128\pi^2} h_t(t)^4 \varphi_c^4(t) \right\}. \end{aligned} \quad (38)$$

Since the expression of the order- \hbar term of the above equation is different from the one we analysed before, we may think at first sight that we would obtain a different result for $m_{\phi(2\text{-loop})}$. In fact we follow the same procedure as before, i.e., first differentiate \tilde{V} by $\varphi_c(t)$, use the relation (17) and eliminate $\mathcal{D}\Omega$ term, evaluate $\partial^2 \tilde{V} / \partial \varphi_c(t)^2$ at $\varphi_c(\tilde{t}_v) = v$ under the minimum condition $\partial \tilde{V} / \partial \varphi_c(t) = 0$ at $\varphi_c(\tilde{t}_v) = v$, and we obtain the following expression:

$$\widetilde{m}_{\phi(2\text{-loop})}^2 = \frac{1}{3}\lambda(\tilde{t}_v)v^2$$

$$\begin{aligned}
& +\hbar v^2 \left\{ \frac{1}{6} \beta_\lambda^{(1)}(\tilde{t}_v) - \frac{1}{6} \lambda(\tilde{t}_v) \left[\frac{\beta_{m^2}^{(1)}(\tilde{t}_v)}{m^2(\tilde{t}_v)} - 2\gamma_\phi^{(1)}(\tilde{t}_v) \right] \right. \\
& \quad \left. + \frac{3}{8\pi^2} h_t(\tilde{t}_v)^4 \right\} \\
& + \mathcal{O}(\hbar^2).
\end{aligned} \tag{39}$$

A logarithmic term of the form $-(3/8\pi^2)\hbar v^2 h_t^4 \ln(h_t^2 \xi^2/2)$ which appeared in Eq.(27) is missing. However, we should note that the definition of t has been altered. Remembering that we have chosen the renormalization scale μ to be M_{SUSY} , we must now evaluate the curvature of EP at the value of \tilde{t}_v , which is determined by

$$\tilde{t}_v = \ln \frac{v}{M_{\text{SUSY}}} + \ln \frac{h_t(\tilde{t}_v)}{\sqrt{2}}. \tag{40}$$

Expanding $\lambda(\tilde{t}_v)$ around t_v , we find

$$\begin{aligned}
\lambda(\tilde{t}_v) & \approx \lambda(t_v) + \hbar \beta_\lambda^{(1)}(t_v) [\tilde{t}_v - t_v] \\
& \approx \lambda(t_v) - \hbar \frac{9}{8\pi^2} h_t^4(t_v) \ln \frac{h_t^2(t_v) \xi^2(t_v)}{2} + \dots \dots \dots
\end{aligned} \tag{41}$$

Then substituting the above expression for $\lambda(\tilde{t}_v)$ into Eq.(39), we will obtain essentially the same result as Eq.(27) except for the h_t^2 - and other non-dominant terms. In fact, we have calculated the mass $\tilde{m}_{\phi(2\text{-loop})}$ with the above choice of t , Eq.(37), and found the difference between the two numerical results for $m_{\phi(2\text{-loop})}$ and $\tilde{m}_{\phi(2\text{-loop})}$ being less than 1GeV for $m_t < 200\text{GeV}$ and $M_{\text{SUSY}} < 10\text{TeV}$.

In conclusion we have examined the mass of the lightest Higgs boson in the MSSM beyond the leading logarithm approximation. We have made use of the EP improved by RGE up to the next-to-leading order. We have found that, contrary to the result of Espinosa and Quirós, the next-to-leading-order corrections to the Higgs mass are non-negligible, adding 3–11GeV (3–9GeV) to the values predicted by the RGE approach in the leading logarithm approximation for the range $100\text{GeV} < m_t < 200\text{GeV}$ and for $M_{\text{SUSY}} = 1\text{TeV}$ ($M_{\text{SUSY}} = 10\text{TeV}$). We also found that the predicted values of m_ϕ are stable under the change of the renormalization parameter t when we use the RGE-improved EP which includes the next-to-leading-order contributions.

Acknowledgements

We would like to thank M. Bando, T.Kugo, N. Maekawa, H. Nakano and Y. Okada for discussions and useful comments. K.S. would like to thank Bill Marciano and Rob Pisarski for discussions and the hospitality extended to him at Brookhaven National Laboratory in the summer of 1993, when part of this work was done.

References

- [1] Y. Okada, M. Yamaguchi, and T. Yanagida, *Prog. Theor. Phys.* **85** (1991) 1; *Phys. Lett.* **B262** (1991) 54.
- [2] J. Ellis, G. Ridolfi, and F. Zwirner, *Phys. Lett.* **B257** (1991) 83; **B262** (1991) 477.
- [3] H. Haber and R. Hempfling, *Phys. Rev. Lett.* **66** (1991) 1815.
- [4] R. Barbieri, M. Frigeni, and F. Caravaglios, *Phys. Lett.* **B258** (1991) 167; R. Barbieri, M. Frigeni, *Phys. Lett.* **B258** (1991) 395.
- [5] A. Yamada, *Phys. Lett.* **B 263** (1991) 233.
- [6] J. R. Espinosa and M. Quirós, *Phys. Lett.* **B266** (1991) 389.
- [7] J. L. Lopez and D. V. Nanopoulos, *Phys. Lett.* **B266** (1991) 397.
- [8] A. Brignole, J. Ellis, G. Ridolfi, and F. Zwirner, *Phys. Lett.* **B271** (1991) 123.
- [9] D. Pierce, A. Papadopoulos and S. Johnson, *Phys. Rev. Lett.* **68** (1992) 3678.
- [10] A. Brignole, *Phys. Lett.* **B281** (1992) 284.
- [11] M. Carena, K. Sasaki and C. E. M. Wagner, *Nucl. Phys.* **B381** (1992) 66.
- [12] J. Gunion and A. Turki, *Phys. Rev.* **D39** (1989) 2701; *Phys. Rev.* **D40** (1989) 2325, 2333.
- [13] M. Berger, *Phys. Rev.* **D41** (1990) 225.
- [14] S. P. Li and M. Sher, *Phys.Lett.* **B140** (1984) 339.
- [15] B. Kastening, *Phys. Lett.* **B283** (1992) 287.
- [16] M. Bando, T. Kugo, N. Maekawa, and H. Nakano, *Phys. Lett.* **B301** (1993) 83.
- [17] C. Ford, D. R. T. Jones, P. W. Stephenson, and M. B. Einhorn, *Nucl. Phys.* **B395** (1993) 17.

- [18] S. Coleman and E. Weinberg, *Phys. Rev.* **D7** (1973) 1888.
- [19] J. Ellis, S. Kelley, and D. V. Nanopoulos, *Phys. Lett.* **B260** (1991) 131.
- [20] P. Langacker and M. Luo, *Phys. Rev.* **D44** (1991) 817.
- [21] U. Amaldi, W. de Boer, and H. Füsrtenu, *Phys. Lett.* **B260** (1991) 240.
- [22] M. Carena, T. E. Clark, C. E. M. Wagner, W. A. Bardeen, and K. Sasaki, *Nucl. Phys.* **B369** (1992) 33.
- [23] M. Lindner, M. Sher, and H. W. Zaglauer, *Phys. Lett.* **B228** (1989) 139.

Figure caption

Fig.1

Values of the Higgs boson mass as a function of m_t , for $M_{\text{SUSY}} = 1\text{TeV}$ and $\cos^2 2\beta = 1$ (upper two lines) and $\cos^2 2\beta = 0$ (lower two lines). The solid and dash-dotted lines denote the next-to-leading-order and the leading-order results, respectively.

Fig.2

The same as in Fig. 1, but considering the case for $M_{\text{SUSY}} = 10\text{TeV}$.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9311366v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9311366v1>